


# LEQVIO<sup>®</sup> Dosing and Administration:

## INJECTION INFORMATION FOR YOUR PRACTICE



### LEQVIO PRODUCT INFORMATION


- Single-dose prefilled syringe of LEQVIO 284 mg/1.5 mL for subcutaneous use<sup>1</sup>
  - 27 gauge 1/2-inch needle<sup>2</sup>
- Sterile, clear and colorless to pale yellow solution<sup>1</sup>
  - This color variation is normal and has no impact on the overall quality of LEQVIO
- Store at room temperature, 68 °F to 77 °F (20 °C to 25 °C)<sup>1</sup>
- Does not require refrigeration

### WHO CAN ADMINISTER LEQVIO?

LEQVIO should be administered by a health care provider.<sup>1\*</sup>

\*Physician, nurse practitioner, physician assistant, or other licensed practitioners.

In-office administration gives you the confidence that your patients received their dose<sup>1</sup>



### HOW IS LEQVIO ADMINISTERED?

LEQVIO is administered in just 2 steps<sup>†</sup>:

#### Step 1: Inspect the syringe

- Check the liquid medicine in the syringe before you give the injection—the solution should be clear and colorless to pale yellow<sup>1</sup>

#### Step 2: Inject the medication

- Inject subcutaneously into the abdomen, upper arm, or thigh<sup>1</sup>
- Injections should not be given in areas of active skin disease or injury, such as sunburns, skin rashes, inflammation, or skin infections<sup>1</sup>

No specific label requirement for post-injection–related safety monitoring.

<sup>†</sup>Clean the injection site with an alcohol swab and use standard aseptic technique when administering injection.

### INDICATION

LEQVIO (inclisiran) injection is indicated as an adjunct to diet and statin therapy for the treatment of adults with primary hyperlipidemia, including heterozygous familial hypercholesterolemia (HeFH), to reduce low-density lipoprotein cholesterol (LDL-C).

### IMPORTANT SAFETY INFORMATION

Adverse reactions in clinical trials ( $\geq 3\%$  of patients treated with LEQVIO and more frequently than placebo) were injection site reaction, arthralgia, and bronchitis.

Please click [here](#) for LEQVIO full Prescribing Information.

## 2 INITIAL DOSES, THEN 2 DOSES A YEAR<sup>1</sup>



INITIAL  
DOSE



ANOTHER AT  
3 MONTHS<sup>1</sup>



### Recommended dose:

One prefilled syringe of  
LEQVIO 284 mg/1.5 mL.<sup>1</sup>

The LEQVIO dosing regimen may integrate seamlessly into a patient's health care routine<sup>1</sup>

## ADDITIONAL CONSIDERATIONS

LEQVIO is administered in combination with maximally tolerated statin therapy.

- **Administration:** Office should schedule the 3-month visit and the following 6-month visit at the time of the initial injection
- **Missed dose:** If a planned dose is missed by less than 3 months, administer LEQVIO and maintain dosing according to the patient's original schedule. If a planned dose is missed by more than 3 months, start a new dosing schedule<sup>1\*</sup>
- **Injection site reactions:**
  - Let patients know that in clinical trials, injection site reactions were predominantly mild or occasionally moderate and did not persist over time<sup>1,3</sup>
  - Advise patients to seek medical attention if injection site irritation persists

\*LEQVIO is administered initially, again at 3 months, and then once every 6 months.

## IMPORTANT SAFETY INFORMATION

Adverse reactions in clinical trials ( $\geq 3\%$  of patients treated with LEQVIO and more frequently than placebo) were injection site reaction, arthralgia, and bronchitis.

Please click [here](#) for LEQVIO full Prescribing Information.

**References:** **1.** Leqvio. Prescribing information. Novartis Pharmaceuticals Corp. **2.** Data on file. LEQVIO Container Closure System. Novartis Pharmaceuticals Corp; 2022. **3.** Data on file. Novartis Pharmaceuticals Corp; 2019.

Licensed from Alnylam Pharmaceuticals, Inc.